



PCT/EP200 4 / 0 0 5 4 9 9



INVESTOR IN PEOPLE

REC'D 0.6 JUL 2004	
WIPO	PCT

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ


I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.


BEST AVAILABLE COPY



Signed *P. McHoney*
Dated 7 May 2004

**PRIORITY
DOCUMENT**

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)



The Patent Office

Cardiff Road
Newport
South Wales
NP10 8QQ

Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

1. Your reference **C4360(C)/RJG**

2. Patent application number **0401810.7**
(The Patent Office will fill this part in)

3. Full name, address and postcode of the or of each applicant (underline all surnames) **UNILEVER PLC
UNILEVER HOUSE, BLACKFRIARS
LONDON, EC4P 4BQ**

Patents ADP number (if you know it) **50426956002**

If the applicant is a corporate body, give the country/state of its incorporation **UNITED KINGDOM**

4. Title of the invention **LIQUID BLEACHING COMPOSITION**

5. Name of your agent (if you have one) **ELLIOTT, Peter William**

"Address for Service" in the United Kingdom to which all correspondence should be sent (including the postcode) **PATENT DEPARTMENT, UNILEVER PLC
COLWORTH HOUSE, SHARNBROOK
BEDFORD, MK44 1LQ**

Patents ADP number (if you know it) **1628003**

6. Priority: Complete this section if you are declaring priority from one or more earlier patent applications, filed in the last 12 months.

Country	Priority application number	Date of filing
GB	0313246.1	9th June 2003

7. Divisionals, etc: Complete this section only if this application is a divisional application or resulted from an entitlement dispute (see note f)

Number of earlier application	Date of filing (day / month / year)

8. Is a Patents Form 7/77 (Statement of inventorship and of right to grant of a patent) required in support of this request?

Answer YES if:

- a) any applicant named in part 3 is not an inventor, or
- b) there is an inventor who is not named as an applicant, or
- c) any named applicant is a corporate body.

Otherwise answer NO (See note d)

YES

Patents Form 1/77

9. Accompanying documents: A patent application must include a description of the invention. Not counting duplicates, please enter the number of pages of each item accompanying this form:

Continuation sheets of this form

Description	16	/
Claim(s)	5	/
Abstract	1	/
Drawing(s)	-	

10. If you are also filing any of the following, state how many against each item.

Priority Documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 7/77)

Request for preliminary examination and search (Patents Form 9/77)

1

Request for substantive examination (Patents Form 10/77)

Any other documents (please specify)

Form 23/77

11. I/We request the grant of a patent on the basis of this application.

Signature(s)



Date: 27/01/04

Sandra Jane EDWARDS, Authorised Signatory

12. Name, daytime telephone number and e-mail address, if any, of person to contact in the United Kingdom

Rob GREEN (01234) 22 2650
rob.green@unilever.com

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 08459 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered YES in part 8, a Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- Part 7 should only be completed when a divisional application is being made under section 15(4), or when an application is being made under section 8(3), 12(6) or 37(4) following an entitlement dispute. By completing part 7 you are requesting that this application takes the same filing date as an earlier UK application. If you want the new application to have the same priority date(s) as the earlier UK application, you should also complete part 6 with the priority details.

LIQUID BLEACHING COMPOSITION

Field of Invention

The present invention provides a liquid bleaching
5 composition.

Background of the Invention

The use of bleaching catalysts for stain removal has been developed over recent years. The recent discovery that some
10 catalysts are capable of bleaching effectively in the absence of an added peroxy source has recently become the focus of some interest, for example: WO9965905; WO0012667; WO0012808; WO0029537, and, WO0060045.

15 The shelf life of a product may be regarded as the period of time over which the product may be stored whilst retaining its required quality. A satisfactory shelf life is in many instances a crucial factor for the success of a commercial product. A product with a short shelf life generally
20 dictates that the product is made in small batches and is rapidly sold to the consumer. It is also a concern to the owners of a brand with a short shelf life that the consumer uses the product within the shelf life otherwise the consumer may be inclined to change to a similar product of
25 another brand. In contrast a similar product with a long shelf life may be made in larger batches, held as stock for a longer period of time and the period of time that a consumer stores the product is not of a great concern to the owners of a particular brand. Despite the shelf life being
30 an important consideration the product must also be active during use.

It is an object of the present invention to provide an air bleaching composition that has improved storage properties whilst being active in use.

5 Summary of Invention

We have found that the storage pH conditions required for longevity of a liquid bleaching composition are in conflict with the pH required for providing good bleaching activity.

- 10 The present invention provides a liquid bleaching composition having a pH between 6 and 7 comprising:
- (a) an organic substance which forms a complex with a transition metal for bleaching a substrate with atmospheric oxygen, the bleaching composition upon addition to an
 - 15 aqueous medium providing an aqueous bleaching medium substantially devoid of a peroxygen bleach or a peroxy-based or peroxy-generating bleach system;
 - (b) a pH changing means; and,
 - (c) the balance carriers and adjunct ingredients to 100
 - 20 wt/wt % of the total bleaching composition, wherein upon dilution of the liquid bleaching composition with water the pH of the liquid bleaching composition enters the range pH 7.5 to pH 9.0.

- 25 The present invention also provides a method of bleaching a textile comprising the steps of:
- (i) diluting from 0.5 to 20 g of a concentrated liquid bleaching composition with 1 litre of water, the
 - concentrated liquid bleaching composition having a pH in the
 - 30 range 6 to 7, the liquid bleaching comprising an organic substance which forms a complex with a transition metal for

bleaching a substrate with atmospheric oxygen, the bleaching composition upon addition to an aqueous medium providing an aqueous bleaching medium substantially devoid of a peroxygen bleach or a peroxy-based or peroxy-generating bleach, the
5 dilution providing an aqueous liquid bleaching composition having a pH in the range from 7.5 to pH 9.0;

- (ii) treating a textile with the aqueous liquid bleaching composition; and,
- (iii) rinsing the textile with water; and,
- 10 (iv) drying the textile.

The term "substantially devoid of a peroxygen bleach or a peroxy-based or peroxy-generating bleach system" should be construed within spirit of the invention. It is preferred
15 that the composition has as low a content of peroxy species present as possible. It is preferred that the bleaching formulation contains less than 1 % wt/wt total concentration of peracid or hydrogen peroxide or source thereof, preferably the bleaching formulation contains less than 0.3
20 % wt/wt total concentration of peracid or hydrogen peroxide or source thereof, most preferably the bleaching composition is devoid of peracid or hydrogen peroxide or source thereof. In addition, it is preferred that the presence of alkyl hydroperoxides is kept to a minimum in a bleaching
25 composition comprising the ligand or complex of the present invention.

The present invention also extends to a commercial package together with instructions for its use.

DETAILED DESCRIPTION OF THE INVENTION

THE BLEACH CATALYST

The bleach catalyst per se may be selected from a wide range of transition metal complexes of organic molecules


5 (ligands). In typical washing compositions the level of the organic substance is such that the in-use level is from 0.05 μM to 50 mM, with preferred in-use levels for domestic laundry operations falling in the range 1 to 100 μM . Higher levels may be desired and applied in industrial textile
10 bleaching processes. A mixture of different catalysts may be employed in the bleaching composition.

Suitable organic molecules (ligands) for forming complexes and complexes thereof are found, for example in:

15 GB 9906474.3; GB 9907714.1; GB 98309168.7, GB 98309169.5; GB 9027415.0 and GB 9907713.3; DE 19755493; EP 999050; WO9534628; EP-A-458379; EP 0909809; United States Patent 4,728,455; WO9839098; WO9839406, WO9748787, WO0029537; WO0052124, and WO0060045 the complexes and organic molecule
20 (ligand) precursors of which are herein incorporated by reference. An example of a preferred catalyst is a transition metal complex of MeN4Py ligand (N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-1-aminoethane).

25 The ligand forms a complex with one or more transition metals, in the latter case for example as a dinuclear complex. Suitable transition metals include for example: manganese in oxidation states II-V, iron II-V, copper I-III, cobalt I-III, titanium II-IV, tungsten IV-VI, vanadium II-V
30 and molybdenum II-VI.

An example of a preferred catalyst is a monomer ligand or transition metal catalyst thereof of a ligand having the formula (I):



5

wherein each R is independently selected from: hydrogen, F, Cl, Br, hydroxyl, C1-C4-alkylo-, -NH-CO-H, -NH-CO-C1-C4-alkyl, -NH₂, -NH-C1-C4-alkyl, and C1-C4-alkyl;

10 R1 and R2 are independently selected from:

C1-C4-alkyl,

C6-C10-aryl, and,

a group containing a heteroatom capable of coordinating to a transition metal, wherein at least one of R1 and R2 is the

15 group containing the heteroatom;

R3 and R4 are independently selected from hydrogen, C1-C8 alkyl, C1-C8-alkyl-O-C1-C8-alkyl, C1-C8-alkyl-O-C6-C10-aryl, C6-C10-aryl, C1-C8-hydroxyalkyl, and -(CH₂)_nC(O)OR₅

wherein R₅ is independently selected from: hydrogen, C1-C4-alkyl, n is from 0 to 4, and mixtures thereof; and,


20 X is selected from C=O, -[C(R₆)₂]_Y- wherein Y is from 0 to 3 each R₆ is independently selected from hydrogen, hydroxyl, C1-C4-alkoxy and C1-C4-alkyl.

25 With regard to the above formula (I) it is also particularly preferred that R1 and R2 may also be independently selected

from: C1 to C22-optionally substituted alkyl, and an optionally substituted tertiary amine of the form -C2-C4-alkyl-NR₇R₈, in which R₇ and R₈ are independently selected from the group consisting of straight chain, branched or cyclo C1-C12 alkyl, benzyl, the -C2-C4-alkyl- of the -C2-C4-alkyl-NR₇R₈ may be substituted by 1 to 4 C1-C2-alkyl, or may form part of a C3 to C6 alkyl ring, and in which R₇ and R₈ may together form a saturated ring containing one or more other heteroatoms.


10

Another preferred class of ligands are macropolycyclic rigid ligands of the formula:



wherein m and n are 0 or integers from 1 to 2, p is an integer from 1 to 6, preferably m and n are both 0 or both 1 (preferably both 1), or m is 0 and n is at least 1; and p is 1; and A is a nonhydrogen moiety preferably having no aromatic content; more particularly each A can vary independently and is preferably selected from methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl, C5-C20 alkyl, and one, but not both, of the A moieties is benzyl, and combinations thereof.

Preferably, the macropolycyclic ligand is of the formula:



wherein "R¹" is independently selected from H, and linear or
 5 branched, substituted or unsubstituted C1-C20 alkyl,
 alkylaryl, alkenyl or alkynyl; and all nitrogen atoms in the
 macropolycyclic rings are coordinated with the transition
 metal.

10 Of the macropolycyclic ligands 5,12-dimethyl-1,5,8,12-
 tetraaza-bicyclo[6.6.2]hexadecane is preferred. This ligand
 is most preferred as its manganese complex [Mn(Bcyclam)Cl₂]
 and may be synthesised according to W09839098.

15 The transition metal complex preferably is of the general
 formula (AI):



20 in which:

M represents a metal selected from Mn(II)-(III)-(IV)-
 (V), Cu(I)-(II)-(III), Fe(II)-(III)-(IV)-(V), Co(I)-(II)-
 (III), Ti(II)-(III)-(IV), V(II)-(III)-(IV)-(V), Mo(II)-
 (III)-(IV)-(V)-(VI) and W(IV)-(V)-(VI), preferably from

25 Fe(II)-(III)-(IV)-(V);

L represents the ligand, preferably N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-1-aminoethane, or its protonated or deprotonated analogue;

5 X represents a coordinating species selected from any mono, bi or tri charged anions and any neutral molecules able to coordinate the metal in a mono, bi or tridentate manner;

Y represents any non-coordinated counter ion;

a represents an integer from 1 to 10;

10 k represents an integer from 1 to 10;

n represents zero or an integer from 1 to 10;

m represents zero or an integer from 1 to 20.

pH Jump Composition

15 The present invention is used as a liquid format where the pH of the commercial product is substantially different to that in use.

20 Sorbitol/borate compositions are known from EP 381 262. In a concentrated solution the borate is complexed with the vicinal diol of the sorbitol; upon dilution of the sorbitol/borate composition the borate is liberated from its interaction with the sorbitol resulting in a pH jump. Other pH Jump formulations are disclosed in US 6,509,308 and US
25 5,484,555.

Balance Carriers and Adjunct Ingredients

These are generally surfactants, builders, foam agents, anti-foam agents, solvents, and enzymes. The use and
30 amounts of these components are such that the bleaching

composition performs depending upon economics, environmental factors and use of the bleaching composition.

The air bleach catalyst may be used in a detergent composition specifically suited for stain bleaching purposes, and this constitutes a second aspect of the invention. To that extent, the composition comprises a surfactant and optionally other conventional detergent ingredients. The invention in its second aspect provides an enzymatic detergent composition which comprises from 0.1 - 50 % by weight, based on the total detergent composition, of one or more surfactants. This surfactant system may in turn comprise 0 - 95 % by weight of one or more anionic surfactants and 5 to 100 % by weight of one or more nonionic surfactants. The surfactant system may additionally contain amphoteric or zwitterionic detergent compounds, but this is not normally desired owing to their relatively high cost. The enzymatic detergent composition according to the invention will generally be used as a dilution in water of about 0.05 to 2%.

In general, the nonionic and anionic surfactants of the surfactant system may be chosen from the surfactants described "Surface Active Agents" Vol. 1, by Schwartz & Perry, Interscience 1949, Vol. 2 by Schwartz, Perry & Berch, Interscience 1958, in the current edition of "McCutcheon's Emulsifiers and Detergents" published by Manufacturing Confectioners Company or in "Tenside-Taschenbuch", H. Stache, 2nd Edn., Carl Hauser Verlag, 1981.

Suitable nonionic detergent compounds which may be used include, in particular, the reaction products of compounds having a hydrophobic group and a reactive hydrogen atom, for example, aliphatic alcohols, acids, amides or alkyl phenols with alkylene oxides, especially ethylene oxide either alone or with propylene oxide. Specific nonionic detergent compounds are C₆-C₂₂ alkyl phenol-ethylene oxide condensates, generally 5 to 25 EO, i.e. 5 to 25 units of ethylene oxide per molecule, and the condensation products of aliphatic C₈-C₁₈ primary or secondary linear or branched alcohols with ethylene oxide, generally 5 to 40 EO.

Suitable anionic detergent compounds which may be used are usually water-soluble alkali metal salts of organic sulphates and sulphonates having alkyl radicals containing from about 8 to about 22 carbon atoms, the term alkyl being used to include the alkyl portion of higher acyl radicals. Examples of suitable synthetic anionic detergent compounds are sodium and potassium alkyl sulphates, especially those obtained by sulphating higher C₈-C₁₈ alcohols, produced for example from tallow or coconut oil, sodium and potassium alkyl C₉-C₂₀ benzene sulphonates, particularly sodium linear secondary alkyl C₁₀-C₁₅ benzene sulphonates; and sodium alkyl glyceryl ether sulphates, especially those ethers of the higher alcohols derived from tallow or coconut oil and synthetic alcohols derived from petroleum. The preferred anionic detergent compounds are sodium C₁₁-C₁₅ alkyl benzene sulphonates and sodium C₁₂-C₁₈ alkyl sulphates. Also applicable are surfactants such as those described in

EP-A-328 177 (Unilever), which show resistance to salting-out, the alkyl polyglycoside surfactants described in EP-A-070 074, and alkyl monoglycosides.

- 5 Preferred surfactant systems are mixtures of anionic with nonionic detergent active materials, in particular the groups and examples of anionic and nonionic surfactants pointed out in EP-A-346 995 (Unilever). Especially preferred is surfactant system that is a mixture of an alkali metal
10 salt of a C_{16} - C_{18} primary alcohol sulphate together with a C_{12} - C_{15} primary alcohol 3-7 EO ethoxylate.

The nonionic detergent is preferably present in amounts greater than 10%, e.g. 25-90% by weight of the surfactant
15 system. Anionic surfactants can be present for example in amounts in the range from about 5% to about 40% by weight of the surfactant system.

One skilled in the art will appreciate that some
20 adventitious peroxy species may be in the composition nevertheless it is most preferred that the bleaching composition of the present invention has less than 1%, preferably less than 0.1%, most preferably less than 0.01%, of a peroxy species present. These adventitious peroxy
25 are predominantly alkyl hydroperoxides formed by autooxidation of the surfactants.

The composition may contain additional enzymes as found in WO 01/00768 A1 page 15, line 25 to page 19, line 29, the
30 contents of which are herein incorporated by reference.

Builders, polymers and other enzymes as optional ingredients may also be present as found in WO0060045.

5 Suitable detergency builders as optional ingredients may also be present as found in WO0034427.

10 In the context of the present invention, bleaching should be understood as relating generally to the decolourisation of stains or of other materials attached to or associated with a substrate. However, it is envisaged that the present invention can be applied where a requirement is the removal and/or neutralisation by an oxidative bleaching reaction of malodours or other undesirable components attached to or otherwise associated with a substrate. Furthermore, in the
15 context of the present invention bleaching is to be understood as being restricted to any bleaching mechanism or process that does not require the presence of light or activation by light.

20 The present invention extends to both isotropic and complex liquid compositions and formulations a brief discussion of which follows. Some isotropic formulations are termed 'micro-emulsion' liquids that are clear and thermodynamically stable over a specified temperature range.
25 The 'micro-emulsion' formulation may be water in oil, or oil in water emulsions. Some liquid formulations are macro-emulsions that are not clear and isotropic. Emulsions are considered meta-stable. Concentrated, clear compositions containing fabric softening actives have been disclosed in
30 WO 98/08924 and WO 98/4799, both Procter & Gamble. Such compositions comprise bio-degradable fabric conditioners.

However, both disclose compositions comprising water miscible solvents that do not form water-in-oil micro-emulsions. Clear fabric conditioning compositions have also been disclosed in EP 730023 (Colgate Palmolive), WO 96/19552

5 (Colgate Palmolive), WO 96/33800 (Witco Co.), WO 97/03170 (Procter & Gamble), WO 97/03172 (Procter & Gamble), WO 97/03169 (Procter & Gamble), US 5492636 (Quest Int.) and US 5427697 (Procter & Gamble). Liquid formulations of the present invention may contain for example; monoethoxy quats; 10 AQAs and bis-AQAs; cationic amides; cationic esters; amino/diamino quats; glucamide; amine oxides; ethoxylated polyethyleneimines; enhancement polymers of the form linear amine based polymers, e.g. bis-hexamethylenetriamine; 15 polyamines e.g. TETA, TEPA or PEI polymers.

It is preferred that the liquid bleaching composition does not contain a pH dependent chromophore, indicator, e.g., phenolphthalein. In addition, it is preferred that the liquid bleaching composition does not contain a pH dependent 20 fluorescent indicator. In both the above cases such should not be present such that a perceptible change is observed to the human eye.

The liquid may be contained within a sachet as found in 25 WO02/068577. The sachet is a container within the context of the present invention.

The liquid composition preferably also contains one or more antioxidants as described in WO02/072747 and WO02072746.

The following is a example of a concentrated liquid formulation that the present invention may be incorporated into by adding the catalyst together with the selected stable perfume components. The commercial liquid
 5 formulation has a pH of 7.

Sodium citrate:	3.2 %
Polypropylene glycol:	4.75 %
LAS-acid:	5.6 %
NI 25 9 EO:	6.6 %
LES (anionic surfactant):	10.5 %
Borax:	2.30 %
Sorbitol:	3.35 %
Alcosperce 725:	0.30 %
Coconut fatty acid:	0.73 %
Monoethanolamine:	0.20 %
Fluorescer:	0.125 %
Enzymes	-
Perfume/dye	-

10 Experimental

The following catalyst was used in the experiments 9,9-dihydroxy-2,4-di-(2-pyridyl)-3-methyl-7-(pyridin-2-ylmethyl)-3,7-diaza-bicyclo[3.3.1]nonane-1,5-dicarboxylate

15 Iron (II) dichloride was prepared as described by Heidi Borzel, Peter Comba, Karl S. Hagen, Yaroslaw D. Lampeka, Achim Lienke, Gerald Linti, Michael Merz, Hans Pritzkow, Lyudmyla V. Tsybal in Inorganica Chimica Acta 337 (2002)

407 - 419. WQ0248301 provides synthetic details of similar compounds. This catalyst was incorporated at a concentration of 0.03 % wt/wt in concentrated liquid composition A containing.

5

Liquid A: 6 % LAS, 6 % sLES 3 EO, 6 % Nonionic 7 EO, 0.016 % Proxel GXL, 3.35 % sorbitol, 2.30 % Borax.10 H₂O, 4.75 % MPG, and sufficient NaOH to bring the pH to 7.

10

Method of Determining Catalyst Stability

The residue bleaching activity of bleaching compositions were determined at 40 °C in a H₂O₂ containing NaH₂PO₄.H₂O pH7 buffer and Acid Blue 45 (CAS No. 2861-02-1) as substrate using the following protocol.

15

Samples of 70 mg liquid were diluted in 10.00 ml MilliQ water. We added 45 µl of this solution to an assay of 230 µL containing 20 mM H₂O₂, 75 µM Acid blue 45 and 54 mM NaH₂PO₄.H₂O pH7 buffer.

20

The changes in absorbance at 600 nm were measured for 5 min at 40 °C using a spectrophotometer. The absolute changes in absorbance were correlated to activities obtained with freshly prepared calibration samples. The measured activities were expressed as µMol/l/. The residual amount is the activity measured on t=0 divided by the activity after storage and expressed in percentages.

25

Table 1: The bleaching catalysts stability expressed as residual amount (%) as function of the pH during storage storage at 37 °C.

pH	1 wk	2 wks	3 wks	4 wks
6	96	82	80	78
6.25	97	84	76	80
6.5	99	91	82	84
6.71	100	88	83	80
7	100	79	71	70
7.25	85	67	56	54
7.52	84	62	54	45
7.75	68	43	38	32
8	62	39	33	25

5

The results in Table 1 clearly show a storage advantage in maintaining the concentrated liquid bleaching composition at a pH at 7 or below. Upon dilution of the concentrated liquid bleaching composition the pH of the solution changes to alkali which activates the bleaching composition.

We claim:

1. A liquid bleaching composition having a pH between 6 and 7 comprising:

5

(a) an organic substance which forms a complex with a transition metal for bleaching a substrate with atmospheric oxygen, the bleaching composition upon addition to an aqueous medium providing an aqueous bleaching medium

10 substantially devoid of a peroxygen bleach or a peroxy-based or peroxy-l-generating bleach system;

(b) a pH changing means; and,

15 (c) the balance carriers and adjunct ingredients to 100 wt/wt % of the total bleaching composition,


wherein upon dilution of the liquid bleaching composition with water the pH of the liquid bleaching composition enters
20 the range pH 7.5 to pH 9.0.

2. A liquid bleaching composition according to claim 1 or 2, wherein upon dilution of the liquid bleaching composition with water the pH of the liquid bleaching composition
25 reaches at least pH 8.0.

3. A liquid bleaching composition according to claim 1, wherein the pH changing means is provided by a borax sorbitol pH jump composition.

4. A liquid bleaching composition according to claim 3, wherein the liquid bleaching composition comprises at least 2 % wt/wt of sorbitol and at least 1 wt/wt % borax.

5. A liquid bleaching composition according to any preceding claim, wherein the organic substance is of formula (I):



10 wherein each R is independently selected from: hydrogen, F, Cl, Br, hydroxyl, C1-C4-alkyl-O-, -NH-CO-H, -NH-CO-C1-C4-alkyl, -NH₂, -NH-C1-C4-alkyl, and C1-C4-alkyl;

R1 and R2 are independently selected from:
C1-C4-alkyl,

15 C6-C10-aryl, and,

a group containing a heteroatom capable of coordinating to a transition metal, wherein at least one of R1 and R2 is the group containing the heteroatom;

20 R3 and R4 are independently selected from hydrogen, C1-C8 alkyl, C1-C8-alkyl-O-C1-C8-alkyl, C1-C8-alkyl-O-C6-C10-aryl, C6-C10-aryl, C1-C8-hydroxyalkyl, and -(CH₂)_nC(O)OR₅

wherein R₅ is independently selected from: hydrogen, C1-C4-alkyl, n is from 0 to 4, and mixtures thereof; and,

X is selected from C=O, -[C(R₆)₂]_Y- wherein Y is from 0 to 3

25 each R₆ is independently selected from hydrogen, hydroxyl, C1-C4-alkoxy and C1-C4-alkyl.

6. A liquid bleaching composition according to any one of claims 1 to 4, wherein the organic substance 5,12-dimethyl-1,5,8,12-tetraaza-bicyclo[6.6.2]hexadecane.

5 7. A method of bleaching a textile comprising the steps of:


(i) diluting from 0.5 to 20 g of a concentrated liquid bleaching composition with 1 litre of water, the concentrated liquid bleaching composition having a pH in the range 6 to 7, the liquid bleaching comprising an organic substance which forms a complex with a transition metal for bleaching a substrate with atmospheric oxygen, the bleaching composition upon addition to an aqueous medium providing an aqueous bleaching medium substantially devoid of a peroxygen bleach or a peroxy-based or peroxy-generating bleach, the dilution providing an aqueous liquid bleaching composition having a pH in the range from 7.5 to pH 9.0;

(iv) treating a textile with the aqueous liquid bleaching composition; and,

20 (v) rinsing the textile with water; and,

(iv) drying the textile.

8. A method according to claim 7, wherein the organic substance is of formula (I):



wherein each R is independently selected from: hydrogen, F, Cl, Br, hydroxyl, C1-C4-alkyl, -NH-CO-H, -NH-CO-C1-C4-alkyl, -NH₂, -NH-C1-C4-alkyl, and C1-C4-alkyl;

R1 and R2 are independently selected from:

5 C1-C4-alkyl,

C6-C10-aryl, and,


a group containing a heteroatom capable of coordinating to a transition metal, wherein at least one of R1 and R2 is the group containing the heteroatom;

10 R3 and R4 are independently selected from hydrogen, C1-C8 alkyl, C1-C8-alkyl-O-C1-C8-alkyl, C1-C8-alkyl-O-C6-C10-aryl, C6-C10-aryl, C1-C8-hydroxyalkyl, and $-(CH_2)_n C(O)OR_5$

wherein R5 is independently selected from: hydrogen, C1-C4-alkyl, n is from 0 to 4, and mixtures thereof; and,

15 X is selected from C=O, $-[C(R_6)_2]_Y-$ wherein Y is from 0 to 3 each R6 is independently selected from hydrogen, hydroxyl, C1-C4-alkoxy and C1-C4-alkyl.

9. A method according to claim 7, wherein the organic
20 substance is of formula:



wherein m and n are 0 or integers from 1 to 2, p is an
25 integer from 1 to 6, preferably m and n are both 0 or both 1

(preferably both 1), or m is 0 and n is at least 1; and p is 1; and A is a nonhydrogen moiety preferably having no aromatic content; more particularly each A can vary independently and is preferably selected from methyl, ethyl, 5 propyl, isopropyl, butyl, isobutyl, tert-butyl, C5-C20 alkyl, and one, but not both, of the A moieties is benzyl, and combinations thereof.

10. A liquid bleaching composition according to claim 9,
10 wherein the organic substance 5,12-dimethyl-1,5,8,12-tetraaza-bicyclo[6.6.2]hexadecane.

Abstract of the Invention

The present invention concerns the preservation of a catalyst in a bleaching composition whilst facilitating activation of the catalyst. The bleaching composition is substantially devoid of a peroxygen bleach or a peroxy-based
5 or peroxy-generating bleach system.

PCT/EP2004/005499



**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.